

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

## (19) World Intellectual Property Organization International Bureau





(43) International Publication Date 12 February 2004 (12.02.2004)

**PCT** 

(10) International Publication Number WO 2004/013276 A3

C12N 15/10

(81) Designated States (national): AU, CA, US.

(51) International Patent Classification7: (21) International Application Number:

PCT/IB2003/003681

(22) International Filing Date: 30 July 2003 (30.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/399;150 30 July 2002 (30.07.2002)

(71) Applicant (for all designated States except US): MOR-PHOSYS IP GMBH [DE/DE]; Lena-Christ-Str. 82152 Martinsried/Planegg (DE).

(71) Applicants and

(72) Inventors: PRASSLER, Josef [DE/DE]; Sandstr. 20, 82110 Munich (DE). STARK, Yvonne [DE/DE]; Hofangerstr. 10, 81735 Munich (DE).

Published:

with international search report

before the expiration of the time limit for amending the claims and to be republished in the event of receipt of

(88) Date of publication of the international search report: 29 December 2004

(15) Information about Correction: **Previous Correction:** see PCT Gazette No. 14/2004 of 1 April 2004, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL TRICISTRONIC VECTORS AND USES THEREFOR

**RBSXbaI** EcoRI HindIII CL VH CH1 phoA L-His6 C MORPH

(57) Abstract: A tricistronic vector (i.e., a vector capable of expressing three exogenous genes, which are not fused together, under the control of one promoter) effectively can encode an immunoglobulin-presenting polypeptide and two immunoglobulin (Ig) polypeptides. The encoded Ig-presenting polypeptide is able to associate with at least one of the Ig polypeptides via co-expressed associating agents. A vector according to the present invention particularly is suited for phage display technology, e.g., when the Ig-presenting polypeptide is a phage coat protein and the Ig polypeptides associate to form a Fab.